



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

to go on training your students to the new terms—some of which you or others will doubtless see reasons for changing—to improve as far as possible, fill up blanks, perhaps get up a small manual in which the new terms shall be practically applied, and have faith that sooner or later the best part of your innovations will find their way into scientific use. The plan is an excellent one, it is a new garment which will fit Science well, if that capricious and fantastic and old-fashioned dressing lady can only be induced to try it on.

Always very truly yours,
O. W. HOLMES.

A CURIOUS EGG.

E. E. BARNARD.

One of my hens of the "Dominico" breed is accountable for the presence to-day of a most remarkable egg, which was found in the hen's nest. This singular object measured about three inches in its longest diameter, a round oval in shape, not like the ordinary egg with a large and a small end. The shell was thin and soft to the touch, resembling the "skin" that is found inside an eggshell. Pressing on one end of the egg, a hard object was felt inside the shell. Opening the egg, by cutting with a sharp knife, two eggs were found, one perfect with a hard shell, slightly smaller than the ordinary egg, the other perfect in every respect, save that it possessed no shell. The egg with the shell was enclosed in the white of the other. These two eggs occupied the two ends of the original shell. Upon opening the one with the hard shell it was found to be perfect. Putting the two eggs in separate cups, the one which had the hard shell was slightly smaller and its yolk of a pale yellow; the yolk of the other was somewhat deeper in color.

Here we have a rare phenomenon; first a large egg with slightly soft shell; inside this two eggs, one perfect in a hard shell, the other without shell but otherwise perfect.

NASHVILLE, Tenn., May 9.

BOOKS RECEIVED.

A MEMORIAL OF JOSEPH HENRY. Published by order of Congress, Washington, 1880.

The present volume presents in a handsome and convenient form the historical facts relating to the career of Professor Joseph Henry, and a record of the various ceremonies and memorial exercises celebrated after his death in honor to his memory.

The memorial exercises at the Capital include addresses by President Garfield, Hannibal Hamlin, Robert E. Withers, Professor Asa Gray, William B. Rogers, General Sherman and others.

The concluding words of President Garfield's address may well be quoted as conveying the general esteem in which Professor Henry was held by all who knew him. "Remembering his great career as a man of science, as a man who served his Government with singular ability and faithfulness, who was loved and venerated by every circle, who blessed with the light of friendship the worthiest and the best, whose life added new lustre to the glory of the human race, we shall be most fortunate, if ever in the future, we see his like again."

NOTES.

RECENT experiments by M. Grehaut, prove that the quantity of carbonic acid exhaled by any one individual of an animal species is about constant. Fifty litres of air passed through the lungs of a dog, 9kg. weight, yielded 2.747 gr. of CO₂. Eight days after the experiment was repeated, and

the CO₂ was 2.810 gr. In man, the same volume of air circulating through the lungs, receives 3.333 gr. of CO₂. Irritations and inflammations of the respiratory mucous membrane (e. g. through inhaling sulphurous acid), considerably decrease the exhalation of CO₂. The gas then tends to accumulate in the blood.

GALVANIC GILDING.—M. Rod gives the composition of a bath to be used at temperatures from 50° to 80° C. It consists of 60 parts crystalline phosphate of soda, 10 parts bisulphate of soda, 1 part cyanide of sodium, 2½ parts chloride of gold, and 1,000 distilled water. In order to prepare the bath the water is divided into three portions of 700, 150, and 150 respectively. The phosphate of soda is dissolved in the first lot, the chloride of gold in the second, and the other ingredients in the third. The two first portions are gradually mixed together, and the third is then slowly added. A platinum plate is used as anode.—*Le Monde de la Science*.

RADIATION THROUGH EBONITE.—Captain Abney exhibited at the Physical Society of London, a number of photographic negatives taken by himself and Colonel Festin by radiation through thin sheets of ebonite. The light from the positive pole of an electric lamp was sent through a thin sheet of ebonite $\frac{1}{4}$ in. thick, and photographs taken showed the radiation to have a low wave-length, from 8,000 to 14,000. The carbon points of the lamp could be photographed through the sheet, and Colonel Festin observed the sun's disc through it. The ebonite showed a grained structure, and different samples of ebonite gave different results, but all gave some result, in course of time at least; old ebonite, like that used in some of Mr. Preece's experiments, scattering the light more than new ebonite. Dr. Moser exhibited the passage of the rays through the ebonite to the audience by means of a galvanometer. Professor Guthrie observed that Captain Abney had proved that light as well as heat traversed the ebonite, and Dr. Coffin stated that compositions of ebonite, apparently the same, might vary considerably.

PHOTOGRAPHIC PHOTOMETRY.—A promising application of photography to precise measurement of phenomena of light has been recently tried by M. Janssen. The method is advantageous in that photography reveals the action of the extremely weak luminous and the ultra-violet rays; but the chief advantage lies in the permanence of the results as against the fugitive nature of ordinary photometric comparisons, which, too, require the simultaneous presence of the two light sources. The various amounts of metallic deposit on the photographic plate cannot well be weighed, so M. Janssen measures by the degree of opacity produced. His photometer consists of a frame with sensitised plate, before which is passed at a known rate of uniform motion a shutter having a slit. If this slit were rectangular, a uniform shade would be produced on the plate; but by making it triangular he obtains a variation of shade, decreasing from the side corresponding to the base of the triangle to that corresponding to the apex. It is further proved that the photographic deposit does not increase as rapidly as the luminous intensity. Now, to compare the sensibility of two plates differently prepared, they have merely to be exposed successively in the frame under like conditions, and the points where they show the same opacity being compared to the points of the triangular slit corresponding to them, the ratio of the apertures at those points expresses the ratio of sensibility. Thus the new gelatinobromide of silver plates are proved to be twenty times as sensitive as the collodion plates prepared by the wet process. Again, to compare two luminous sources, they are made to act successively on two similar plates in the photometer, and the points of equal shade in the plates indicate, as before, the relation sought. M. Janssen has compared the light of the sun and some stars on these principles, preparing from the former "solar scales" (with uniform degradation of shade), under exactly determined conditions as to sensitive layer, time of solar action, height of the sun, etc. Circular images of stars are obtained by placing a photographic plate a little out of focus in the telescope, and a series of these, got with different times of exposure, are compared with the scales obtained from sunlight. M. Janssen will shortly make known some of his results.